

Masahiro Nakamura · Takayuki Ohgushi

Positive and negative effects of leaf shelters on herbivorous insects: linking multiple herbivore species on a willow

Received: 13 August 2002 / Accepted: 22 April 2003 / Published online: 24 May 2003
© Springer-Verlag 2003

Abstract We experimentally examined the effects on other herbivorous insects of leaf shelters constructed by lepidopteran larvae on a willow, *Salix miyabeana*. Several insect species occupied the vacant leaf shelters. Our experiment using artificial leaf shelters showed that the number of aphids increased with the number of artificial leaf shelters on a shoot, as did the numbers of three ant species (*Camponotus japonicus*, *Lasius hayashi*, and *Myrmica jessensis*) that entered leaf shelters to collect aphid honeydew. To determine the ant-mediated effect of leaf shelters on herbivorous insects that do not use leaf shelters, we transferred newly hatched larvae of a common leaf beetle, *Plagioderia versicolora*, to the leaves of shoots with and without artificial leaf shelters. One day after the transfer, larval survival rate was significantly lower on shoots with shelters than on those without shelters, and shoots with shelters had significantly more ants than did shoots without shelters. Our field experiments demonstrated clearly that shelter-making lepidopteran larvae increased the abundance of both aphids and ants and decreased the survival rate of leaf beetle larvae, probably because the larvae were removed by ants that were attracted to the leaf shelters by the aphid colonies.

Keywords Ant-aphid mutualism · Habitat resources · Shelter-makers · Secondary users · Shelter-mediated interactions

Introduction

The prevalence of interspecific interactions among herbivorous insects has long been discounted in insect community ecology (Hairston et al. 1960; Lawton and Strong 1981). Recently, however, there has been increasing evidence that herbivorous insects sharing a host plant often do affect one other considerably by changing the quality or quantity of the host plant (Faeth 1986, 1987; Karban and Myers 1989; Hunter and West 1990; Damman 1993; Denno et al. 1995; Ohgushi 1997). Herbivory by one species often induces morphological, phenological, and chemical changes in the host plant that, in turn, alter the availability of resources for other species that feed at different times or on different parts of the plant. Such indirect plant-mediated interactions may commonly occur between species that are separated by time or space (Faeth 1986, 1987; Harrison and Karban 1986; Gange and Brown 1989; Moran and Whitham 1990; Masters and Brown 1992; Masters et al. 1993; Ohgushi 1997).

In the study of plant-mediated interactions, much attention has been paid to changes in the quality or quantity of host plants following herbivory, because plants have mainly been considered simply as food for herbivorous insects (Faeth 1986, 1987; Karban and Myers 1989). However, plants provide not only food but also microhabitats that protect many insect species against natural enemies or environmental stress (Strong et al. 1984; Hunter and West 1990).

Several studies have reported that leaf shelters, such as leaf rolls and leaf galls made by herbivorous insects, are later used by other arthropods as microhabitats (Carroll and Kearby 1978; Carroll et al. 1979; Akimoto 1981; Hajek and Dahlsten 1986; Cappuccino 1993; Cappuccino and Martin 1994; Kudo 1994; Larsson et al. 1997; Alper 1998; Martinsen et al. 2000). For example, Akimoto (1981) reported that the aphid *Eriosoma yangi* inhabited leaf-roll galls on elm trees made by other *Eriosoma* species. Since herbivorous insects commonly make shelters (Martinsen et al. 2000), shelter-mediated inter-

M. Nakamura (✉) · T. Ohgushi
Institute of Low Temperature Science, Hokkaido University,
060–0819 Sapporo, Japan
e-mail: masahiro@ecology.kyoto-u.ac.jp
Tel.: +81-77-5498019
Fax: +81-77-5498231

Present address:

M. Nakamura · T. Ohgushi, Center for Ecological Research,
Kyoto University, Kamitanakami, Otsu, 520–2113 Shiga, Japan

actions between arthropods that make shelters and those that do not may occur frequently on terrestrial plants.

In Hokkaido, northern Japan, numerous species of lepidopteran larvae construct leaf shelters such as leaf folds, rolls, sandwiches, and ties on a species of willow tree, *Salix miyabeana* (Seemen). Several species of arthropods later occupy the vacant leaf shelters after the lepidopteran larvae have emerged or made new shelters. In particular, we have frequently observed an aphid, *Chaitophorous saliniger* (Shinji), in leaf shelters. After these aphids have colonized the leaf shelters, ants visit the shelters, probably to obtain aphid honeydew. Therefore, we predict that leaf shelters on *S. miyabeana* decrease the larval survivorship of a common leaf beetle, *Plagioderia versicolora* (Laicharting) because attending ants remove the larvae.

The aims of this study were to determine whether leaf shelters affect the abundance of the aphid and its attending ants and whether the presence of leaf shelters subsequently affects the larval survivorship of *P. versicolora*, which does not use leaf shelters. We determined the number and species of leaf-shelter occupants, tested the effect of artificial leaf shelters on the abundance of aphids and attending ants, and examined the subsequent effects on the survivorship of leaf beetle larvae.

Materials and methods

Study site and host plant

This study was performed in 1999 on the banks of the Ishikari River (43°N, 141°E) in Hokkaido, the northern island of Japan. Six willow species (*Salix miyabeana* Seemen, *S. sachalinensis* Fr. Schmidt, *S. integra* Thunberg, *S. hultenii* Floderus, *S. subfragilis* Anders, and *S. pet-susu* Kimura) were sympatric at the study site. Field observations and experiments were conducted on *S. miyabeana*, the most abundant willow species in the study area (Ishihara et al. 1999), which supported a high density of leaf shelters built by lepidopteran larvae.

Shelter-makers and secondary users

A number of lepidopteran larvae, mainly in the Tortricidae and Pyralidae families, make leaf shelters on *S. miyabeana* by folding or rolling a single leaf or by tying two or three leaves together with silk. The larvae either develop in the leaf shelters until pupation or move on and make new shelters. Vacant shelters are often occupied by other arthropods (secondary users).

Leaf beetle

The leaf beetle *Plagioderia versicolora* Laicharting is a dominant specialist herbivorous insect that feeds on *Salix* and *Populus* leaves (Kimoto and Takizawa 1994). Overwintering adults begin to emerge in late May; females lay egg clutches on the undersurface of leaves from early June to late August; and larvae pass through three instars and then pupate (Ishihara et al. 1999).

Survey of secondary users

To determine the species and numbers of secondary users of leaf shelters, we randomly selected 10–15 trees on which we sampled leaf shelters from mid-May until late September. In mid-May and late September, we sampled only 50 leaf shelters, because relatively few were found, but otherwise we sampled 100 leaf

shelters three times a month. While sampling, we lifted leaf shelters carefully to avoid displacing secondary users and immediately packed the shelters in small vinyl bags. The bags were kept in a laboratory freezer at –30°C. We classified the leaf shelters into the following categories: Containing lepidopteran larvae, *Chaitophorous saliniger*, *C. saliniger* with ants, other arthropods, and without secondary users.

Effects of leaf shelters on abundance of aphids and ants

To determine whether leaf shelters affect the abundance of *C. saliniger* and its attending ants, we conducted a field experiment involving artificial leaf shelters. Each artificial leaf shelter was constructed by rolling a single leaf and maintaining the rolled shape by binding the ends with two plastic tubes (6 mm diameter, 10 mm length). On 12 July 1999, we randomly selected 23 trees that were 2–3 m in height. On each tree, we selected four current-year shoots of similar size. On each shoot, we set 0, 1, 3, or 6 artificial leaf shelters, with each shoot receiving a different number. Two weeks later, the shoots were individually packed in vinyl bags and frozen at –30°C. We then counted the number of *C. saliniger* aphids and their attendant ants on each shoot. To determine the relationships between leaf shelters and aphids and between leaf shelters and ants, we used a linear regression of log ($n+1$)-transformed numbers of aphids and ants. To determine the relationship between aphids and ants, we used a linear regression of the raw data.

Effects of leaf shelters on survival of leaf beetle larvae

We conducted a larval transfer experiment to determine how leaf shelters affect the survival rate of *P. versicolora* larvae, which may be removed by ants attending aphid colonies in the leaf shelters. On 15 July 1999, we randomly selected 11 trees that were 2–3 m in height. On each tree, we selected two current-year shoots of similar size. On each shoot, we set four artificial leaf shelters on one shoot (the shelter shoot) and none on the other (the non-shelter shoot). Two weeks later, newly hatched beetle larvae [11.3 ± 2.2 larvae (mean \pm 1 SD)] were transferred to a leaf on each shoot. During the first 6 h after the transfer, we counted the number of ants on each shoot every hour. One day later, we counted the number of remaining beetle larvae on each shoot to estimate the larval survival rate. We used the Mann-Whitney *U* test to examine differences in larval survival rates and in the cumulative number of ants on shelter shoots and non-shelter shoots.

Results

Secondary users

Of the 7,927 arthropods in the sampled leaf shelters, 94.4% were aphids (*Chaitophorous saliniger*), 5.0% were ants (*Camponotus japonicus*, *Lasius hayashi*, and *Myrmica jessensis*) that were attending aphid colonies in the shelters, and 0.6% were other arthropods.

Species composition in leaf shelters changed with season (Fig. 1). The proportion of leaf shelters housing lepidopteran larvae decreased consistently from mid-May to mid-July because of adult emergence. In late July, second-generation lepidopteran larvae began to construct leaf shelters. The number of lepidopteran larvae was highest in mid-August and declined thereafter. In contrast, *C. saliniger* began to colonize vacant leaf shelters in mid-June; in mid-July and mid-September, the aphid had colonized approximately 75% of all leaf shelters.




Fig. 1 Proportion of leaf shelters made on *Salix miyabeana* containing different types of occupants. Shelters were classified into five categories: (1) lepidopteran larvae, (2) aphids (*Chaitophorus saliniger*), (3) aphids with ants, (4) other arthropods, and (5) no occupants




Fig. 2A, B Effect of artificial leaf shelters on the abundance of aphids (*C. saliniger*) and its attending ants. Relationship between number of artificial leaf shelters and **A** aphids and **B** ants

Effects of leaf shelters on aphid and ant abundance

The number of aphids and associated ants significantly increased with increasing number of artificial leaf shelters on a shoot (aphid: $n=92$, $r=0.708$, $P<0.0001$; ants; $n=92$, $r=0.724$, $P<0.0001$) (Fig. 2). The number of ants increased significantly with the number of aphids ($n=92$, $r=0.790$, $P<0.0001$) (Fig. 3).




Fig. 3 Relationship between number of aphids (*C. saliniger*) and attending ants




Fig. 4A, B Ant-mediated effect of artificial leaf shelters on the survival rate of transferred larvae of the leaf beetle, *Plagioder a versicolora*, near leaf shelters. **A** proportion of leaf beetles surviving one day after transfer to shelter shoots and non-shelter shoots, and **B** cumulative number of ants on shelter shoots and non-shelter shoots over the first six hours after larval transfer. Means \pm 1 SE are presented

Effects of leaf shelters on survival of leaf beetle larvae

The presence of leaf shelters negatively affected the survival rate of beetle larvae near leaf shelters. One day after larval transfer, the larval survival rate was significantly lower on shelter shoots than on non-shelter shoots ($n=11$, $U=19.5$, $P=0.0065$) (Fig. 4A). In contrast to the few ants observed on non-shelter shoots, ants visited artificial leaf shelters continuously. Thus, the cumulative number of ants during the first 6 h was over 74 times higher than the number on non-shelter shoots ($n=11$, $U=8.5$, $P=0.0003$) (Fig. 4B).

Discussion

Effects of leaf shelters on other arthropods

This study showed that the leaf shelters constructed by lepidopteran larvae were used as microhabitats by several insect species, the most abundant of which was the aphid, *Chaitophorous saliniger*. Similarly, recent studies have shown that shelter-makers interact positively with secondary users by providing leaf shelters, which increase microhabitat availability (Cappuccino 1993; Cappuccino and Martin 1994; Larsson et al. 1997; Martinsen et al. 2000; Fukui 2001).

We also found that leaf shelters positively affected the abundance of an aphid and attending ants that collected honeydew. This increase in ant numbers subsequently negatively affected the larval survival rate of a leaf beetle, *Plagioderia versicolora*, near leaf shelters.

Positive effects of leaf shelters on aphid and ant abundance

Several authors have suggested that most interactions among herbivorous insects are indirect and that such interactions can be positive as well as negative (Damman 1993; Denno et al. 1995). Our experiments clearly illustrated that the lepidopteran larvae positively affected aphid abundance by providing leaf shelters. This might be due to increased survival of the aphids and a subsequent increase in reproduction. Living in a leaf shelter has several benefits: refuge from natural enemies (Fowler and MacGarvin 1985; Damman 1987; Atlegrim 1989; Cappuccino 1993; Eubanks et al. 1997), protection against adverse microclimatic conditions (Henson 1958; Hunter and Willmer 1989; Larsson et al. 1997), and acquisition of high-quality food (Lewis 1979; Dussourd and Eisner 1987; Sandberg and Berenbaum 1989; Sagers 1992; Fukui et al. 2002).

Ants are often strongly associated with aphid colonies, because honeydew is a stationary and renewable food resource (Way 1963; Carroll and Janzen 1973; Buckley 1987). In this study, three ant species were found with *C. saliniger*, the aphid that colonized leaf shelters, and the abundance of these ants correlated highly with the number of the aphid. We conclude, therefore, that the ants were probably gathering aphid honeydew.

Some authors have argued that bottom-up regulation by host-plant characteristics is the primary factor controlling herbivorous insect population dynamics (Hunter and Price 1992; Ohgushi 1992; Harrison and Cappuccino 1995). This argument focuses on the quantity and quality of host plants as food for herbivores (Ohgushi and Sawada 1985; Schultz 1988). We showed, however, that an increase in leaf shelters increased aphid abundance. Therefore, the aphid population may be limited by the abundance of leaf shelters. This conclusion implies that, in addition to food availability, microhabitat availability on host plants is also an important factor in determining the population dynamics of secondary users.

Negative effects of leaf shelters on survival of leaf beetle larvae

Leaf shelters may negatively affect herbivorous insects near leaf shelters if predators are attracted to the shelters (Martinsen et al. 2000; Fukui 2001). In our larval transfer experiment, attending ants were often observed patrolling outside leaf shelters. These ants frequently encountered leaf beetle larvae and removed them from the trees. One day after larval transfer, survival was much lower on shelter shoots, where the attending ants probably removed the transferred beetle larvae. Many authors have reported negative effects of the ant-aphid association on other herbivorous insects (Messina 1981; Fritz 1983; Fowler and MacGarvin 1985; Ito and Higashi 1991; Floate and Whitham 1994; Wimp and Whitham 2001), because attending ants remove herbivorous competitors of the aphid food resources (Way 1963; Buckley 1987). In the field, we commonly found one or two leaf shelters on a shoot during the experimental period (mid-July), which is somewhat lower than the density used in the larval transfer experiment.

In conclusion, our field experiments suggest that shelter-makers provide critical microhabitats to other herbivorous insects. These positive shelter-mediated interactions among herbivorous insects on terrestrial plants may be much more common and widespread than previously thought. This is because insects including Lepidoptera, weevils, sawflies, and aphids construct leaf shelters on a wide variety of plants, from trees and shrubs to herbs and even ferns (Martinsen et al. 2000). On the other hand, this study revealed that by attracting predators leaf shelters can also negatively affect herbivorous insects that do not use the leaf shelters. Future studies need to focus on both direct and indirect effects of leaf shelters on herbivorous insects, to establish the prevalence of positive and negative effects.

Acknowledgements We thank M. Kondoh, M. Walker, and T. Craig for their valuable comments on earlier drafts of this manuscript. We also thank T. Hayashi for help in the field. This study was partially supported by a Grant-in-Aid for General Scientific Research to T.O. (09NP1501) from the Ministry of Education, Science, Sports and Culture, Japan, and by the MEXT Grant-in-Aid for the 21st Century COE Program (A2 to Kyoto University).

References

- Akimoto S (1981) Gall formation by *Eriosoma fundatrices* and gall parasitism in *Eriosoma yangi* (Homoptera, Pemphigidae). *Kontyu* 49:426–436
- Alper J (1998) Ecosystem engineers shape habitats for other species. *Science* 280:1195–1196
- Atlegrim O (1989) Exclusion of birds from bilberry stands: impact on insect larval density and damage to the bilberry. *Oecologia* 79:136–139
- Buckley RC (1987) Interactions involving plants, homoptera, and ants. *Annu Rev Ecol Syst* 18:111–135
- Cappuccino N (1993) Mutual use of leaf-shelters by lepidopteran larvae on paper birch. *Ecol Entomol* 18:287–292

- Cappuccino N, Martin M (1994) Eliminating early-season leaf-tiers of paper birch reduces abundance of mid-summer species. *Ecol Entomol* 19:399–401
- Carroll CR, Janzen DH (1973) Ecology of foraging by ants. *Annu Rev Ecol Syst* 4:231–257
- Carroll MR, Kearby WH (1978) Microlepidopterous oak leaf-tiers (Lepidoptera: Gelechioidea) in central Missouri. *J Kans Entomol Soc* 51:457–471
- Carroll MR, Wooster MT, Kearby WH, Allen DC (1979) Biological observations on three oak leaf-tiers: *Psilocorsis quercicella*, *P. reflexella* and *P. cryptolechiella* in Massachusetts and Missouri. *Ann Entomol Soc Am* 72:441–447
- Damman H (1987) Leaf quality and enemy avoidance by the larvae of a pyralid moth. *Ecology* 68:88–97
- Damman H (1993) Patterns of interaction among herbivore species. In: Stamp NE, Casey TM (eds) *Caterpillars: ecological and evolutionary constraints on foraging*. Chapman and Hall, New York, pp 132–169
- Denno RF, McClure MS, Ott JR (1995) Interspecific interactions in phytophagous insects: competition reexamined and resurrected. *Annu Rev Entomol* 40:297–331
- Dussourd DE, Eisner T (1987) Vein-cutting behavior: insect counterploy to the latex defense of plants. *Science* 237:898–901
- Eubanks MD, Nesci KA, Petersen MK, Lui Z, Sanchez HB (1997) The exploitation of an ant-defended host plant by a shelter-building herbivore. *Oecologia* 109:454–460
- Faeth SH (1986) Indirect interactions between temporally separated herbivores mediated by the host plant. *Ecology* 67:479–494
- Faeth SH (1987) Community structure and folivorous insect outbreaks: the roles of vertical and horizontal interactions. In: Barbosa P, Schultz JC (eds) *Insect outbreaks*. Academic Press, San Diego, pp 135–171
- Floate KD, Whitham TG (1994) Aphid-ant interaction reduces chrysomelid herbivory in a cottonwood hybrid zone. *Oecologia* 97:215–221
- Fowler SV, MacGarvin M (1985) The impact of hairy wood ants, *Formica lugubris*, on the guild structure of herbivorous insects on birch, *Betula pubescens*. *J Anim Ecol* 54:847–855
- Fritz RS (1983) Ant protection of a host plant's defoliator: consequence of an ant-membracid mutualism. *Ecology* 64:789–797
- Fukui A (2001) Indirect interactions mediated by leaf shelters in animal-plant communities. *Popul Ecol* 43:31–40
- Fukui A, Murakami M, Konno K, Nakamura M, Ohgushi T (2002) A leaf-rolling caterpillar improves leaf quality. *Entomol Sci* 5:263–266
- Gange AC, Brown VK (1989) Effects of root herbivory by an insect on a foliar-feeding species, mediated through changes in the host plant. *Oecologia* 81:38–42
- Hairston NG, Smith FE, Slobodkin LB (1960) Community structure, population control, and competition. *Am Nat* 44:421–425
- Hajek AE, Dahlsten DL (1986) Coexistence of three species of leaf-feeding aphids (Homoptera) on *Betula pendula*. *Oecologia* 68:380–386
- Harrison S, Cappuccino N (1995) Using density-manipulation experiments to study population regulation. In: Cappuccino N, Price PW (eds) *Population dynamics: new approaches and synthesis*. Academic Press, San Diego, pp 131–147
- Harrison S, Karban R (1986) Effects of an early-season folivorous moth on the success of a later-season species, mediated by a change in the quality of the shared host, *Lupinus arboreus* Sims. *Oecologia* 69:354–359
- Henson WR (1958) The effects of radiation on the habitat temperatures of some poplar-inhabiting insects. *Can J Zool* 36:463–478
- Hunter MD, Price PW (1992) Playing chutes and ladders: heterogeneity and the relative roles of bottom-up and top-down forces in natural communities. *Ecology* 73:724–732
- Hunter MD, West C (1990) Variation in the effects of spring defoliation on the late season phytophagous insects of *Quercus robur*. In: Watt AD, Leather SR, Hunter MD, Kidd NAC (eds) *Population dynamics of forest insects*. Intercept, Edinburgh, Scotland, pp 123–135
- Hunter MD, Willmer PG (1989) The potential for interspecific competition between two abundant defoliators on oak: leaf damage and habitat quality. *Ecol Entomol* 14:267–277.
- Ishihara M, Hayashi T, Ohgushi T (1999) Life cycle of the willow leaf beetle, *Plagioderia versicolora* (Coleoptera: Chrysomelidae) in Ishikari (Hokkaido, Japan). *Entomol Sci* 2:57–60
- Ito F, Higashi S (1991) An indirect mutualism between oaks and wood ants via aphids. *J Anim Ecol* 60:463–470
- Karban R, Myers JH (1989) Induced plant responses to herbivory. *Annu Rev Ecol Syst* 20:331–348
- Kimoto S, Takizawa H (1994) Leaf beetles (Chrysomelidae) of Japan (in Japanese). Tokai University Press, Tokyo
- Kudo S (1994) Observations on lepidopteran leaf-shelters as molting refuges for the stink bug *Elasmucha putoni* (Heteroptera: Acanthosomatidae). *Psyche* 101:183–186
- Larsson S, Hågström H, Denno RF (1997) Preference for protected feeding site by larvae of the willow-feeding leaf beetle *Galerucella lineola*. *Ecol Entomol* 22:445–452
- Lawton JH, Strong DR (1981) Community patterns and competition in folivorous insects. *Am Nat* 118:317–338
- Lewis AC (1979) Feeding preference for diseased and wilted sunflower in the grasshopper, *Melanoplus differentialis*. *Entomol Exp Appl* 26:202–207
- Martinsen GD, Floate KD, Waltz AM, Wimp GM, Whitham TG (2000) Positive interactions between leafrollers and other arthropods enhance biodiversity on hybrid cottonwoods. *Oecologia* 123:82–89
- Masters GJ, Brown VK (1992) Plant-mediated interactions between two spatially separated insects. *Funct Ecol* 6:175–179
- Masters GJ, Brown VK, Gange AC (1993) Plant mediated interactions between above- and below-ground insect herbivores. *Oikos* 66:148–151
- Messina FJ (1981) Plant protection as a consequence of an ant-membracid mutualism: interactions on goldenrod (*Solidago* sp.). *Ecology* 62:1433–1440
- Moran NA, Whitham TG (1990) Interspecific competition between root-feeding and leaf-galling aphids mediated by host-plant resistance. *Ecology* 71:1050–1058
- Ohgushi T (1992) Resource limitation on insect herbivore population. In: Hunter MD, Ohgushi T, Price PW (eds) *Effect of resource distribution on animal-plant interactions*. Academic Press, San Diego, pp 199–241
- Ohgushi T (1997) Plant-mediated interactions between herbivorous insects. In: Abe T, Levin SA, Higashi M (eds) *Biodiversity: an ecological perspective*. Springer, Berlin Heidelberg New York, pp 115–130
- Ohgushi T, Sawada H (1985) Population equilibrium with respect to available food resource and its behavioural basis in an herbivorous lady beetle *Henosepilachna niponica*. *J Anim Ecol* 54:781–796
- Sagers CL (1992) Manipulation of host plant quality: herbivores keep leaves in the dark. *Funct Ecol* 6:741–743
- Sandberg SL, Berenbaum MR (1989) Leaf-tying by tortricid larvae as an adaptation for feeding on phototoxic *Hypericum perforatum*. *J Chem Ecol* 15:875–885
- Schultz JC (1988) Many factors influence the evolution of herbivore diets, but plant chemistry is central. *Ecology* 69:896–897
- Strong DR, Lawton JH, Southwood TRE (1984) *Insects on plants: community patterns and mechanisms*. Blackwell, Oxford
- Way MJ (1963) Mutualism between ants and honeydew-producing homoptera. *Annu Rev Entomol* 8:307–344
- Wimp GM, Whitham TG (2001) Biodiversity consequences of predation and host plant hybridization on an aphid-ant mutualism. *Ecology* 82:440–452